



## Fight-It 32™



CLEANER • DISINFECTANT • DETERGENT • DEODORIZER

### RECOMMENDATIONS:

- FOOD PROCESSING FACILITIES
- GENERAL CLEANING
- DISINFECTING, DEODORIZING
- SANITIZE NON-FOOD CONTACT SURFACES
- DISINFECT FOOD SERVICE ESTABLISHMENT FOOD CONTACT SURFACES
- USE IN: hospitals and healthcare settings, home health care, hospices, laundry rooms, nursing homes, catering facilities, banquet halls, institutional kitchens and/or bathroom.

### KEY FEATURES/BENEFITS:

#### \*Virucide

**Fungicide** (against pathogenic fungi)

**Mildewstat** (on hard non-porous inanimate surfaces)

#### ACTIVE INGREDIENT:

Octyl decyl dimethyl ammonium chloride .....	0.814%
Dioctyl dimethyl ammonium chloride .....	0.407%
Didecyl dimethyl ammonium chloride .....	0.407%
Alkyl (40% <sub>C<sub>12</sub></sub> , 50% <sub>C<sub>14</sub></sub> , 10% <sub>C<sub>16</sub></sub> )	
dimethyl benzyl ammonium chloride .....	1.085%

### PHYSICAL PROPERTIES:

Physical State: Liquid  
 Appearance: clear, orange liquid  
 Odour: orange  
 pH: 5.5-7.5  
 Specific Gravity: 1.00  
 Solubility in Water: Very Soluble  
 Viscosity: As Water  
 Percent Volatile by Volume: 94% by weight

### DIRECTIONS:

**DISINFECTION:** To disinfect inanimate, hard non-porous surfaces add 32 mL of this product per Litre of water. Apply solution with a mop, cloth, sponge or hand-pump trigger sprayer so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes, then remove excess liquid. For heavily soiled areas, a pre-cleaning step is required. Prepare a fresh solution for each use.



42366  
4/4L Concentrated Formula

42367  
1/20L Concentrated Formula

**TO DISINFECT TOILET BOWLS:** Remove gross filth or soils from surfaces with bowl brush. Add 32 mL of product to the bowl water. Brush or swab the bowl completely using a scrub brush or toilet mop making sure to get under the rim. Let stand for 10 minutes and flush.

**DEODORIZATION:** To deodorize, apply this product as indicated under the heading DISINFECTION.

**TO DISINFECT FOOD PROCESSING ESTABLISHMENTS:** To disinfect food processing premises, floors, walls, storage area and equipment, add 32 mL per litre of water. Apply solution with a mop, cloth, sponge or hand pump trigger sprayer so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes, then remove excess liquid. Before using this product, food products and packaging materials must be removed from the area and carefully protected. After use, all surfaces in the area must be rinsed with potable water.

**BACTERICIDAL ACTIVITY:** At the 32 mL per Litre dilution, this product demonstrates effective activity against the organisms:

- Bordetella bronchiseptica
- Corynebacterium ammoniagenes
- Enterobacter aerogenes
- Enterobacter cloacae
- Enterobacter cloacae (clinical isolate)
- Enterococcus faecalis (clinical isolate)
- Enterococcus faecalis
- Escherichia coli
- Enterococcus faecalis (Vancomycin resistant)
- Escherichia coli (clinical isolate)
- Fusobacterium necrophorum
- Lactobacillus casei subs. rhamnosus
- Klebsiella pneumoniae subsp. pneumoniae
- Listeria monocytogenes

Pasteurella multocida  
Proteus mirabilis ATCC 25933  
Proteus mirabilis ATCC 9921  
Proteus vulgaris  
Pseudomonas acroginosa  
Salmonella choleraesuis  
Salmonella choleraesuis subsp. choleraesuis serotype typhimurium  
Salmonella choleraesuis subsp. choleraesuis serotype paratyphi B  
Salmonella choleraesuis subsp. choleraesuis serotype pullorum  
Salmonella choleraesuis subsp. choleraesuis serotype typhi  
Serratia marcescens  
Shigella sonnei  
Shigella flexneri Type 2b  
Shigella dysenteriae  
Staphylococcus aureus  
Staphylococcus aureus subsp. aureus  
Staphylococcus aureus (Methicillin resistant)  
Staphylococcus aureus (clinical isolate)  
Staphylococcus epidermidis (clinical isolate)  
Staphylococcus epidermidis  
Streptococcus pyogenes Group A  
Staphylococcus aureus (Vancomycin Intermediate)  
Streptococcus pyogenes (clinical flesh eating strain - Bird M3)  
Xanthomonas maltophilia (clinical isolate)

**FUNGICIDAL ACTIVITY:** At the 32 mL per Litre dilution, this product demonstrates effective fungicidal activity against the pathogenic fungi Trichophyton mentagrophytes and Candida albicans.

EFFICACY TESTS HAVE DEMONSTRATED THAT THIS PRODUCT IS AN EFFECTIVE FUNGICIDE IN WATER UP TO 200 PPM HARDNESS (AS CaCO<sub>3</sub>) IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).

**MILDEWSTAT:** To control mold and mildew (such as Aspergillus niger) and the odours they cause on pre-cleaned, hard, non-porous surfaces add 32 mL per litre of water. Apply solution with a cloth, mop or sponge making sure to wet all surfaces completely. Let air dry. Prepare a fresh solution for each use. Repeat application at weekly intervals or when mildew growth reappears.

**\*VIRUCIDAL ACTIVITY:** This product when used on environmental, inanimate hard non-porous surfaces at 32 mL per Litre of water exhibits effective virucidal activity against:

Herpes Simplex Type 1 (causative agent for fever blisters)  
Herpes Simplex Type 2 (genital disease)  
Influenza A2/Hong Kong  
Vaccinia  
Pseudorabies  
Bovine Rhinotracheitis  
Feline Leukemia  
Feline Picornavirus  
Canine Distemper virus  
Human Immunodeficiency Virus Type 1 and Type 2 (HIV-1 and HIV-2)

**KILLS HIV-1 AND HIV-2 ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential transmission of Human Immunodeficiency Virus Type 1 and Type 2 (HIV-1 and HIV-2).

Special Instructions for Cleaning and Decontamination against HIV-1 and HIV-2 of Surfaces Soiled with Blood or Body Fluids.

Personal Protection: When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks, or eye coverings.

Cleaning Procedures: Blood or body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

Contact Time/Dilution: At a use dilution of 1 part per 32 parts water FIGHT-IT 32™ is effective against HIV-1 and HIV-2 in the presence of 5% blood serum with a 10 minute contact time.

Disposal of Infectious Materials: Blood or other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

EFFICACY TESTS HAVE DEMONSTRATED THAT THIS PRODUCT IS AN EFFECTIVE BACTERICIDE AND VIRUCIDE IN WATER UP TO 400 PPM HARDNESS (AS CaCO<sub>3</sub>) IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).

#### PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. DANGER - CORROSIVE TO EYES - SKIN IRRITANT. Causes eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. If swallowed, drink large quantities of water. Avoid alcohol. Call a physician or poison control centre immediately. Take container, label or product name and Registration Number with you when you seek medical attention.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

DISPOSAL: Rinse the emptied container thoroughly. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

#### SIZES:

**42366:** 4/4L Concentrated Formula  
**42367:** 1/20L Concentrated Formula  
**42368:** 1/205L Concentrated Formula

**SDS AVAILABLE UPON REQUEST #0379**